

## Computer Aided Geometric Design

### Curves

### Bezier Curve

**Instructor: Dr. Prashant K. Jain**  
 Professor ME Discipline  
 PDPM Indian Institute of Information Technology,  
 Design and Manufacturing Jabalpur, Jabalpur, INDIA  
 Resources: [web.iitdmj.ac.in/~pkjain/](http://web.iitdmj.ac.in/~pkjain/)  
 Email: [pkjain@iitdmj.ac.in](mailto:pkjain@iitdmj.ac.in), [pkjain2006@gmail.com](mailto:pkjain2006@gmail.com)  
<http://www.iitdmj.ac.in/Faculty/pkjain.html>  
<http://in.linkedin.com/in/pkjain2006>  
<https://www.facebook.com/pkjain2006>



**PDPM**  
**INDIAN INSTITUTE OF INFORMATION TECHNOLOGY,**  
**DESIGN AND MANUFACTURING JABALPUR**  
 (An Institute of National Importance (INI) established by MHRD, Govt. of INDIA)

### Bezier Curve

Mathematically a parametric Bezier curve is defined by

$$P(t) = \sum_{i=0}^n B_i J_{n,i}(t)$$

where the Bezier or Bernstein basis or blending function is

$$J_{n,i}(t) = \binom{n}{i} t^i (1-t)^{n-i}$$

With

$$\binom{n}{i} = \frac{n!}{i!(n-i)!}$$


$J_{n,i}(t)$  is the  $i$ -th  $n$ th order Bernstein basis function, here  $n$  is the degree of the defining Bernstein basis function.

Thus degree of the polynomial curve segment, is one less than the number of points in the defining Bezier polygon.

September 16, 2024 Dr. Prashant K. Jain (IITDMJ)

### Blending functions

Blending functions for several values of  $n$   
 Notice the symmetry of the functions



(a) three polygon points  $n=2$ ; (b) four polygon points,  $n=3$ ;  
 (c) five polygon points,  $n=4$ ; (d) six polygon points,  $n=5$ .


September 16, 2024 Dr. Prashant K. Jain (IITDMJ)

For example, each of the four blending functions shown in Figure for  $n=3$  is a cubic. The maximum value of each blending function occurs at  $t=i/n$ .

$$J_{n,i}(t) = \binom{n}{i} \frac{i^i (n-i)^{n-i}}{n^n}$$

For example, for a cubic  $n=3$ , The maximum values for  $J_{3,1}$  and  $J_{3,2}$  occur at  $1/3$  and  $2/3$ , respectively, with values

$$J_{3,1}\left(\frac{1}{3}\right) = \frac{4}{9}$$

$$J_{3,2}\left(\frac{2}{3}\right) = \frac{4}{9}$$


September 16, 2024 Dr. Prashant K. Jain (IITDMJ)

The derivatives of the basis function are obtained by formally differentiating equation  $J_{n,i}(t) = \binom{n}{i} t^i (1-t)^{n-i}$ . Specifically

$$J'_{n,i}(t) = \binom{n}{i} \{ i t^{i-1} (1-t)^{n-i} - (n-i) t^i (1-t)^{n-i-1} \} = 0$$

$$i t^{i-1} (1-t)^{n-i} - (n-i) t^i (1-t)^{n-i-1} = 0$$

$$i t^{i-1} (1-t)^{n-i} = (n-i) t^i (1-t)^{n-i-1}$$

$$\frac{i t^i}{t} (1-t)^{n-i} = (n-i) t^i \frac{(1-t)^{n-i}}{(1-t)}$$

$$\frac{i}{t} = \frac{(n-i)}{(1-t)}$$

$$\frac{(1-t)}{t} = \frac{(n-i)}{i}$$

$$\frac{1}{t} - 1 = \frac{n-i}{i} - 1$$

$$\frac{1}{t} = \frac{n}{i}$$

September 16, 2024 Dr. Prashant K. Jain (IITDMJ)

Examining eqns. for the first point on the curve i.e. at  $t=0$  shows that

$$J_{n,0}(0) = \frac{n! (0)^i (1-0)^{n-i}}{n! (n-i)!} = 1 \quad i=0$$

$$J_{n,i}(0) = \frac{n! (0)^i (1-0)^{n-i}}{n! (n-i)!} = 0 \quad i \neq 0$$

Thus  $P(0) = B_0 J_{n,0}(0) = B_0$

This shows that the first point on the Bezier curve and on its defining polygon are coincident.

Similarly for the last point on the curve, i.e. at  $t=1$

$$J_{n,n}(1) = \frac{n! (1)^i (0)^{n-i}}{n! (1)!} = 1 \quad i=n$$

$$J_{n,i}(1) = \frac{n! (1)^i (1-1)^{n-i}}{n! (n-i)!} = 0 \quad i \neq n$$

Thus  $P(1) = B_n J_{n,n}(1) = B_n$

This shows that the last point on the Bezier curve and the last point on its defining polygon are coincident.

September 16, 2024 Dr. Prashant K. Jain (IITDMJ)

### Derivatives Bezier Curves

Although it is not necessary to numerically specify the tangent vectors at the ends of an individual Bezier curve, maintaining slope and curvature continuity when joining Bezier curves, determining surface normal for lighting or numerical control tool path calculations, or local curvature for smoothness or fairness calculations requires a knowledge of both first and second derivatives of a Bezier curve.

Recalling equation, the first derivative of a Bezier curve is

$$P'(t) = \sum_{i=0}^n B_i J'_{n,i}(t)$$

Second derivative is given by

$$P''(t) = \sum_{i=0}^n B_i J''_{n,i}(t)$$

September 16, 2024

Dr. Prashant K. Jain (IIITDM)

9

### Derivatives Bezier Curves

The derivatives of the basis function are obtained by formally differentiating

equation  $J_{n,i}(t) = \binom{n}{i} t^i (1-t)^{n-i}$ . Specifically

$$\begin{aligned} J'_{n,i}(t) &= \binom{n}{i} \{i t^{i-1} (1-t)^{n-i} - (n-i) t^i (1-t)^{n-i-1}\} \\ &= \binom{n}{i} t^i (1-t)^{n-i} \left\{ \frac{i}{t} - \frac{(n-i)}{(1-t)} \right\} \\ &= \frac{(i-nt)}{t(1-t)} J_{n,i}(t) \end{aligned}$$

Similarly the second derivative is

$$J''_{n,i}(t) = \left\{ \frac{(i-nt)^2 - nt^2 - i(i-2t)}{t^2(1-t)^2} \right\} J_{n,i}(t)$$

September 16, 2024

Dr. Prashant K. Jain (IIITDM)

10

### Derivatives Bezier Curves

At the beginning and the ends of a Bezier curve i.e. at  $t=0$  and  $t=1$  numerical evaluation of above equations creates difficulties.

An alternate evaluation for the  $r$ th derivative at  $t=0$  is given by

$$P^r(0) = \frac{n!}{(n-r)!} \sum_{i=0}^r (-1)^{r-i} \binom{r}{i} B_i$$

At  $t=1$  by

$$P^r(1) = \frac{n!}{(n-r)!} \sum_{i=0}^r (-1)^i \binom{r}{i} B_{n-i}$$

Thus the first derivatives at the ends are

$$P'(0) = n(B_1 - B_0)$$

And

$$P'(1) = n(B_n - B_{n-1})$$

This illustrates that the tangent vector for a Bezier curve at the initial and final points has the same direction as the initial and final polygon spans.

September 16, 2024

Dr. Prashant K. Jain (IIITDM)

11

### Derivatives Bezier Curves

Similarly the second derivatives at the ends are

$$P''(0) = n(n-1)(B_0 - 2B_1 + B_2)$$

And

$$P''(1) = n(n-1)(B_n - 2B_{n-1} + B_{n-2})$$

Thus the second derivative of the Bezier curve at the initial and final points depends on the two nearest polygon spans i.e. on the nearest three polygon vertices.

September 16, 2024

Dr. Prashant K. Jain (IIITDM)

12

### Example on Derivatives

Consider the four points Bezier polygon

$$P(t) = B_0 J_{3,0}(t) + B_1 J_{3,1}(t) + B_2 J_{3,2}(t) + B_3 J_{3,3}(t)$$

Hence first derivative is

$$P'(t) = B_0 J'_{3,0}(t) + B_1 J'_{3,1}(t) + B_2 J'_{3,2}(t) + B_3 J'_{3,3}(t)$$

Differentiating the basis functions directly yields

$$J_{3,0}(t) = t^0(1-t)^3 \rightarrow J'_{3,0}(t) = -3(1-t)^2$$

$$J_{3,1}(t) = 3t(1-t)^2 \rightarrow J'_{3,1}(t) = 3(1-t)^2 - 6t(1-t)$$

$$J_{3,2}(t) = 3t^2(1-t) \rightarrow J'_{3,2}(t) = 6t(1-t) - 3t^2$$

$$J_{3,3}(t) = t^3 \rightarrow J'_{3,3}(t) = 3t^2$$

September 16, 2024

Dr. Prashant K. Jain (IIITDM)

13

### Example on Derivatives

Consider the four points Bezier polygon

$$P(t) = B_0 J_{3,0}(t) + B_1 J_{3,1}(t) + B_2 J_{3,2}(t) + B_3 J_{3,3}(t)$$

Hence first derivative is

$$P'(t) = B_0 J'_{3,0}(t) + B_1 J'_{3,1}(t) + B_2 J'_{3,2}(t) + B_3 J'_{3,3}(t)$$

Second derivative is

$$P''(t) = B_0 J''_{3,0}(t) + B_1 J''_{3,1}(t) + B_2 J''_{3,2}(t) + B_3 J''_{3,3}(t)$$

Differentiating the basis functions directly yields

$$J_{3,0}(t) = t^0(1-t)^3 \rightarrow J'_{3,0}(t) = -3(1-t)^2 \rightarrow J''_{3,0}(t) = 6(1-t)$$

$$J_{3,1}(t) = 3t(1-t)^2 \rightarrow J'_{3,1}(t) = 3(1-t)^2 - 6t(1-t) \rightarrow J''_{3,1}(t) = -6(2-3t)$$

$$J_{3,2}(t) = 3t^2(1-t) \rightarrow J'_{3,2}(t) = 6t(1-t) - 3t^2 \rightarrow J''_{3,2}(t) = 6(1-3t)$$

$$J_{3,3}(t) = t^3 \rightarrow J'_{3,3}(t) = 3t^2 \rightarrow J''_{3,3}(t) = 6t$$

September 16, 2024

Dr. Prashant K. Jain (IIITDM)

14

**Example on Derivatives**

Evaluating the results at  $t=0$  yields

$$J'_{3,0}(0) = -3, \quad J'_{3,1}(0) = 3, \quad J'_{3,2}(0) = 0, \quad J'_{3,3}(0) = 0$$

Substituting yields

$$P'(0) = -3B_0 + 3B_1 = 3(-B_0 + B_1)$$

Thus the direction of the tangent vector at the beginning of the curve is the same as that of the first polygon span

At the end of the curve,  $t=1$  and

$$J'_{3,0}(1) = 0, \quad J'_{3,1}(1) = 0, \quad J'_{3,2}(1) = -3, \quad J'_{3,3}(1) = 3$$

Substituting yields

$$P'(1) = -3B_2 + 3B_3 = 3(-B_2 + B_3)$$

Thus, the direction of the tangent vector at the end of the curve is the same as that of the last polygon span.

September 16, 2024

Dr. Prashant K. Jain (IITDM)

15

**Example on Derivatives**

Evaluating the results at  $t=0$  yields

$$J''_{3,0}(0) = 6, \quad J''_{3,1}(0) = -12, \quad J''_{3,2}(0) = 6, \quad J''_{3,3}(0) = 0$$

Substituting yields

$$P''(0) = 6B_0 - 12B_1 + 6B_2 = 6(B_0 - 2B_1 + B_2)$$

Thus second derivative or curvature of Bezier curve at start point depends on three nearest polygon points or two nearest polygon spans.

At the end of the curve,  $t=1$  and

$$J''_{3,0}(1) = 0, \quad J''_{3,1}(1) = 6, \quad J''_{3,2}(1) = -12, \quad J''_{3,3}(1) = 6$$

Substituting yields

$$P''(1) = 6B_1 - 12B_2 + 6B_3 = 6(B_1 - 2B_2 + B_3)$$

Thus second derivative or curvature of Bezier curve at end point depends on three nearest polygon points or two nearest polygon spans.

September 16, 2024

Dr. Prashant K. Jain (IITDM)

16

**Example on Derivatives**

Evaluating the results at  $t=0$  yields

$$J'_{3,0}(0) = -3, \quad J'_{3,1}(0) = 3, \quad J'_{3,2}(0) = 0, \quad J'_{3,3}(0) = 0$$

Substituting yields

$$P'(0) = -3P_0 + 3P_1 = 3(-P_0 + P_1)$$

Thus the direction of the tangent vector at the beginning of the curve is the same as that of the first polygon span

At the end of the curve,  $t=1$  and

$$J'_{3,0}(1) = 0, \quad J'_{3,1}(1) = 0, \quad J'_{3,2}(1) = -3, \quad J'_{3,3}(1) = 3$$

Substituting yields

$$P'(1) = -3P_2 + 3P_3 = 3(-P_2 + P_3)$$

Thus, the direction of the tangent vector at the end of the curve is the same as that of the last polygon span.

September 16, 2024

Dr. Prashant K. Jain (IITDM)

17

The basis functions given above along with equations can be used to evaluate the derivations along the curve. Specifically, the first derivatives are

$$J'_{3,0}(t) = \frac{(0-3t)}{t(1-t)}(1-t)^3 = -3(1-t)^2$$

$$J'_{3,1}(t) = \frac{(1-3t)}{t(1-t)}(3t)(1-t)^2 = 3(1-3t)(1-t) = 3(1-4t+3t^2)$$

$$J'_{3,2}(t) = \frac{(2-3t)}{t(1-t)}(3t^2)(1-t) = 3t(2-3t)$$

$$J'_{3,3}(t) = \frac{3(1-t)}{t(1-t)}t^3 = (3t^2)$$

Notice that there is no difficulty in evaluating these results at either  $t=0$  or  $t=1$ . Substituting into eqn. yields at first derivative at any point on the curve.

For example, at  $t=1/2$ ,

$$\begin{aligned} P'\left(\frac{1}{2}\right) &= -3\left(1-\frac{1}{2}\right)^2 B_0 + 3\left(1-\frac{3}{2}\right)\left(1-\frac{1}{2}\right) B_1 + \left(\frac{3}{2}\right)\left(2-\frac{3}{2}\right) B_2 + \frac{3}{4} B_3 \\ &= -\frac{3}{4} B_0 - \frac{3}{4} B_1 + \frac{3}{4} B_2 + \frac{3}{4} B_3 = -\frac{3}{4} (B_0 + B_1 - B_2 - B_3) \end{aligned}$$

September 16, 2024

Dr. Prashant K. Jain (IITDM)

18

Similarly, the second derivatives are

$$J''_{3,0}(t) = \frac{\{(-3t)^2 - 3t^2\}}{t^2(1-t)^2}(1-t)^3 = 6(1-t)$$

$$\begin{aligned} J''_{3,1}(t) &= \frac{\{(1-3t)^2 - 3t^2 - (1-2t)\}}{t^2(1-t)^2}(3t)(1-t)^2 \\ &= -6(2-3t) \end{aligned}$$

$$\begin{aligned} J''_{3,2}(t) &= \frac{\{(2-3t)^2 - 3t^2 - 2(1-2t)\}}{t^2(1-t)^2}3t^2(1-t) \\ &= 6(1-3t) \end{aligned}$$

$$J''_{3,3}(t) = \frac{\{(3-3t)^2 - 3t^2 - 3(1-2t)\}}{t^2(1-t)^2}t^3 = 6t$$

For  $t=1/2$ , these results yield

$$\begin{aligned} P''\left(\frac{1}{2}\right) &= 6\left(1-\frac{1}{2}\right) B_0 - 6\left(2-\frac{3}{2}\right) B_1 + 6\left(1-\frac{3}{2}\right) B_2 + 3B_3 \\ &= 3B_0 - 3B_1 - 3B_2 + 3B_3 = 3(B_0 - B_1 - B_2 + B_3) \end{aligned}$$

September 16, 2024

Dr. Prashant K. Jain (IITDM)

20

**Continuity conditions**

If one Bezier curve  $P(t)$  of degree  $n$  is defined by vertices  $B_i$  and an adjacent Bezier curve  $Q(s)$  of degree  $m$  by vertices  $C_i$  then first derivative continuity at the joint between the curves is given by

$$P'(1) = gQ'(0)$$

Where  $g$  is a scalar.

$$n(B_n - B_{n-1}) = m(C_1 - C_0)$$

$$(C_1 - C_0) = \frac{n}{m}(B_n - B_{n-1})$$

Since positional continuity is implied at the joint,  $C_0 = B_n$  and

$$C_1 = \frac{n}{m}(B_n - B_{n-1}) + B_n$$

Thus the tangent vector directions at the joint are the same if the three vertices

$B_{n-1}$ ,  $B_n = C_0$  and  $C_1$  are collinear, i.e.  $B_n$  need only lie somewhere on the line between  $B_{n-1}$  and  $C_1$ .

September 16, 2024

Dr. Prashant K. Jain (IITDM)

22

If both direction and magnitude of the tangent vectors at the joint are to be equal then for equal degree curves ( $n=m$ ),  $B_n = C_0$  must be at midpoint of the line joining  $B_{n-1}$  and  $C_1$  i.e.

$$(C_1 - C_0) = B_n - B_{n-1} = C_0 - B_{n-1}$$

$$C_1 + B_{n-1} = 2C_0 = 2B_n$$

Figure illustrates this for  $n = m = 3$  i.e. For two cubic Bezier curves.

September 16, 2024 Dr. Prashant K. Jain (IIITDM) 23

Second derivative continuity between adjacent Bezier curves is given by

$$m(m-1)(C_0 - 2C_1 + C_2) = n(n-1)(B_{n-2} - 2B_{n-1} + B_n)$$

Using the conditions for  $C_0$  and  $C_1$  continuity at the joint given above yields

$$C_2 = \frac{n(n-1)}{m(m-1)} B_{n-2} - 2 \left\{ \frac{n(n-1)}{m(m-1)} + \frac{n}{m} \right\} B_{n-1} + \left\{ 1 + 2 \frac{n}{m} + \frac{n(n-1)}{m(m-1)} \right\} B_n$$

For cubic Bezier curve

$$C_2 = B_{n-2} - 4(B_{n-1} - B_n)$$

Second derivative continuity between quartic Bezier curve

September 16, 2024 Dr. Prashant K. Jain (IIITDM) 24

### Additional flexibility in Bezier Curves

- Degree raising
- Sub division

September 16, 2024 Dr. Prashant K. Jain (IIITDM) 25

### Additional flexibility in Bezier curve

#### Sub division

Additional flexibility can also be obtained by subdividing a Bezier curve into two new Bezier curves that, combined are identical with the original curve. A Bezier curve can be divided at any parameter value in the range 0 to 1. The simplest choice at the midpoint i.e.  $t=1/2$ .

Here the results for midpoint subdivision are derived for the special case of cubic Bezier curves.

A cubic Bezier curve is given by

$$p(t) = (1-t)^3 B_0 + 3t(1-t)^2 B_1 + 3t^2(1-t) B_2 + t^3 B_3 \quad 0 \leq t \leq 1$$

September 16, 2024 Dr. Prashant K. Jain (IIITDM) 26

### Additional flexibility in Bezier curve

#### Sub division

September 16, 2024 Dr. Prashant K. Jain (IIITDM) 27

### Sub division of Bezier Curve

With defining polygon given by  $B_0, B_1, B_2, B_3$ . The polygon  $C_0, C_1, C_2, C_3$  then defining the Bezier curve  $Q(u)$ ,  $0 \leq u \leq 1$  corresponding to the first half of the original curve i.e.  $p(t) = 0 \leq t \leq 1/2$  is required.

Similarly, the polygon  $D_0, D_1, D_2, D_3$  that defines the Bezier curve  $R(v)$ ,  $0 \leq v \leq 1$  corresponding to the second half of the original curve;  $p(t) = 1/2 \leq t \leq 1$  is also required.

The new defining polygon vertices  $C_i$  and  $D_i$  are obtained by equating the position and tangent vectors

at  $u = 0, t = 0$ ;  $u = 1, t = 1/2$  and  $v = 0, t = 1/2$ ;  $v = 1, t = 1$ ;

Using Bezier definitions

We obtain these equations

$$C_0 = B_0$$

$$3(C_1 - C_0) = \frac{3}{2}(B_1 - B_0)$$

$$3(C_3 - C_2) = \frac{3}{8}(B_3 + B_2 + B_1 - B_0)$$

$$C_3 = \frac{1}{8}(B_3 + 3B_2 + 3B_1 + B_0)$$

$$P(t) = \sum_{i=0}^n B_i J_{n,i}(t)$$

$$P'(t) = \sum_{i=0}^n B_i J'_{n,i}(t)$$

September 16, 2024 Dr. Prashant K. Jain (IIITDM) 28

### Sub division of Bezier Curve

Solution of these equations gives

$$C_0 = B_0$$

$$C_1 = \frac{1}{2}(B_1 + B_0)$$

$$C_2 = \frac{1}{4}(B_2 + 2B_1 + B_0)$$

$$C_3 = \frac{1}{8}(B_3 + 3B_2 + 3B_1 + B_0)$$

Results generalize to

$$C_i = \sum_{j=0}^i \binom{i}{j} \frac{B_j}{2^i} \quad i = 0, 1, \dots, n$$

$$D_i = \sum_{j=i}^n \binom{n-i}{n-j} \frac{B_j}{2^{n-i}} \quad i = 0, 1, \dots, n$$

Applied successfully, the defining polygons converge to the Bezier curve itself


September 16, 2024

Dr. Prashant K. Jain (IIITDM)

30

### Sub division of Bezier Curve

Define a Bezier curve with four polygon vertices  $B_0[0 \ 2 \ 4]$ ,  $B_1[4 \ 6 \ 4]$ ,  $B_2[6 \ 1 \ 4]$  and  $B_3[10 \ 5 \ 4]$ , then Subdivide this curve in two separate Bezier curves at  $t=0.4$ . Plot two new control polygons.



| First Control Polygon |       |   |
|-----------------------|-------|---|
| 0                     | 2     | 4 |
| 1.6                   | 3.6   | 4 |
| 2.88                  | 3.76  | 4 |
| 4.096                 | 3.632 | 4 |

| Second Control Polygon |       |   |
|------------------------|-------|---|
| 4.096                  | 3.632 | 4 |
| 5.92                   | 3.44  | 4 |
| 7.6                    | 2.6   | 4 |
| 10                     | 5     | 4 |

September 16, 2024

Dr. Prashant K. Jain (IIITDM)

31

### de Casteljau's Algorithm

#### Curve Subdivision

##### ◆ Cascading lerps

$$p_{01} = (1-t)p_0 + tp_1$$

$$p_{12} = (1-t)p_1 + tp_2$$

$$p_{23} = (1-t)p_2 + tp_3$$

$$p_{012} = (1-t)p_{01} + tp_{12}$$

$$p_{123} = (1-t)p_{12} + tp_{23}$$


$$p_{0123} = (1-t)p_{012} + tp_{123}$$

##### ◆ Subdivides curve at $p_{0123}$

$$\blacksquare p_0 \ p_{01} \ p_{012} \ p_{0123}$$

$$\blacksquare p_{0123} \ p_{123} \ p_{23} \ p_3$$

##### ◆ Repeated subdivision converges to curve




September 16, 2024

Dr. Prashant K. Jain (IIITDM)

32

### Trimming of Bezier Curve

Now complete relationship between two sets of control polygons



$$q_i = p_i$$

$$q_j = p_j$$

$$q_i' = \frac{u_j - u_i}{v_j - v_i} p_i'$$

$$q_j' = \frac{u_j - u_i}{v_j - v_i} p_j'$$

➤ Tangent vector magnitudes must change to accommodate a change in the range of the parametric variable and are simply scaled by the ratio of the ranges of parametric variable.

➤ This preserves the direction of tangent vectors and shape of the curve.


September 16, 2024

Dr. Prashant K. Jain (IIITDM)

33

### Trimming of Bezier Curve

Define a Bezier curve with four polygon vertices  $B_0[4 \ 4]$ ,  $B_1[2 \ 8]$ ,  $B_2[6 \ 8]$  and  $B_3[8 \ 4]$ , then Trim this curve in a smaller new Bezier curves at  $t=0.3$  to  $t=0.7$ . Plot new control polygon and curve.



September 16, 2024

Dr. Prashant K. Jain (IIITDM)

34

### Additional flexibility in Bezier curve

#### Degree Raising

If a Bezier curve with additional flexibility is required, the degree of the defining polynomial can be increased by increasing the number of defining polygon points.

For every point on a Bezier curve with  $n$  defining polygon vertices  $B_0 \dots B_n$  the same point on a new Bezier curve with  $n+1$  defining polygon vertices is given by  $B_0^* \dots B_{n+1}^*$

$$P(t) = \sum_{i=0}^n B_i J_{n,i}(t) = \sum_{i=0}^{n+1} B_i^* J_{n+1,i}(t)$$

$$\text{Where } B_0^* = B_0$$

$$B_i^* = \alpha_i B_{i-1} + (1 - \alpha_i) B_i$$





$$\alpha_i = \frac{i}{n+1} \quad i = 1, \dots, n$$

$$B_{n+1}^* = B_n$$

September 16, 2024

Dr. Prashant K. Jain (IIITDM)

35



# THANK YOU

**Dr. Prashant K. Jain**  
Professor (ME Design)

**PDPM IIITDM Jabalpur**  
INDIAN INSTITUTE OF INFORMATION TECHNOLOGY, DESIGN & MANUFACTURING JABALPUR  
(An Institute of National Importance, established by GOI, Govt. of India)

Ph: +91-767-2794415  
Cell: +91-94268-03010  
Email: prajain@iiitdm.ac.in  
prajain2008 @prajain2008

Chairman, Rajaratnam Fund, IITD, Kharanpur  
Jabalpur: 482 005, MP, India  
http://www.iiitdm.ac.in